

Distribution of millipedes along an altitudinal gradient in the south of Lake Teletskoye, Altai Mts, Russia (Diplopoda)

Julia S. Nefedieva¹, Pavel S. Nefediev²,
Miroslava B. Sakhnevich³, Yuri V. Dyachkov²

1 Barnaul Branch of OJSC “GIPRODORNII”, Papanintsev street 105, Barnaul, 656000, Russia **2** Department of Ecology, Biochemistry and Biotechnology, Altai State University, Lenina avenue 61, Barnaul, 656049, Russia **3** Altai State Nature Biosphere Reserve, Naberezhnyi lane 1, Gorno-Altaysk, 649000, Russia

Corresponding author: Julia S. Nefedieva (j.nefedieva@mail.ru)

Academic editor: Ivan H. Tuf | Received 28 October 2014 | Accepted 26 May 2015 | Published 30 June 2015

<http://zoobank.org/9A4EB171-797E-415A-8872-0F0182099AFA>

Citation: Nefedieva JS, Nefediev PS, Sakhnevich MB, Dyachkov YV (2015) Distribution of millipedes along an altitudinal gradient in the south of Lake Teletskoye, Altai Mts, Russia (Diplopoda). In: Tuf IH, Tajovský K (Eds) Proceedings of the 16th International Congress of Myriapodology, Olomouc, Czech Republic. ZooKeys 510: 141–161. doi: 10.3897/zookeys.510.8855

Abstract

The distribution of millipedes along an altitudinal gradient in the south of Lake Teletskoye, Altai, Russia based on new samples from the Kyga Profile sites, as well as on partly published and freshly revised material (Mikhaljova et al. 2007, 2008, 2014, Nefedieva and Nefediev 2008, Nefediev and Nefedieva 2013, Nefedieva et al. 2014), is established. The millipede diversity is estimated to be at least 15 species and subspecies from 10 genera, 6 families and three orders. The bulk of species diversity is confined both to low- and mid-mountain chern taiga forests and high-mountain shrub tundras, whereas the highest numbers, reaching up to 130 ind./m², is shown in subalpine *Pinus sibirica* sparse growths. Based on clustering studied localities on species diversity similarity two groups of sites are defined: low-mountain sites and subalpine sparse growths of *P. sibirica* ones.

Keywords

Millipedes, Diplopoda, altitudinal gradient, Lake Teletskoye, Altai, Siberia

Introduction

This paper continues ecological researches on the Altai millipede fauna in the south of Lake Teletskoye, Russian Altai (Nefedieva and Nefediev 2008, Nefediev and Nefedieva 2013). Some faunistic records of *Sibiriuulus altaicus* Gulička, 1972 and specimens of the family Diplomaragnidae from the study localities have been made earlier (Mikhaljova et al. 2007, 2008, 2014). A brief historical account of Altai millipede fauna research can be obtained from the publication of Mikhaljova et al. (2008) and Nefediev and Nefedieva (2013).

Since 1998 the Altai State Nature Biosphere Reserve and a buffer zone around Lake Teletskoye are inscribed as one of three separate areas of UNESCO World Natural Heritage Site under the name of «Golden Mountains of Altai». Lake Teletskoye, being the deepest and the largest body of freshwater in southwest Siberia, exerts a great warming influence on local climate, in its southern part especially. Situated at a height of 435 m above sea level, the lake lies between the mountain ridges of Altyntu and Korbu, and the Chulyshman river highlands in the south.

The aim of our present paper is to explore the distribution of millipedes along an altitudinal gradient in the south of the Teletskoye Lake in the Kyga Biogeocenosis Profile.

Material and methods

Material was collected by the first and the second authors of this article in August 2005 using hand sampling from the litter and standard technique of soil sampling (Ghilarov 1987): 8 soil samples per each studied numbered site, sample area $\frac{1}{4}$ m², depth 10 cm. The total amount of studied millipedes is 968 specimens.

The Kyga Biogeocenosis Profile was laid in 1959–1961 in the territory of watershed of the Kyga and Bayas rivers in the south of the Teletskoye Lake near the cordon of Chiri, Ulagan District, Republic of Altai, Siberia, Russia. The profile encompasses 22 numbered sites at different altitudes, and comprising relic *Pinus sibirica* forests from the Tertiary period. Its length is about 12 km, and its altitude ranges from 443 to 2267 m a.s.l. It is begun at the mouth of the Kyga river and ends on the top of the Malaya Koliushhta mountain. The vertical vegetation zonation here is characterized by the presence of forest and high-mountain belts. There are widespread dark coniferous forests with *Betula pendula* and *Populus tremula*, also called as chern taiga, and sparse growths of *P. sibirica* in the former belt, whereas alpine meadows do not occur almost at all in the latter. Above the timberline at a height of 2100 m above sea level, all hilltops are occupied by shrub, moss-lichen and rocky tundras.

All sites we collected are listed and described below according to the following standard: site number (**bold**): GPS (WGS84) position, altitude, habitat, sampling date, sampling methods.

- 1:** 51°20'47,3"N, 87°51'14,2"E, 443 m a.s.l., *Pinus sylvestris* and *Betula pendula* forest with *Larix sibirica*, *Abies sibirica* and *P. sibirica*, 12.08.2005, soil sampling.

- 2:** 51°20'29,3"N, 87°51'40,0"E, 494 m a.s.l., *A. sibirica* and *P. sibirica* forest with ferns, 12.08.2005, soil sampling.
- 4:** 51°19'53,3"N, 87°51'78,0"E, 675 m a.s.l., *A. sibirica* forest with *P. sibirica* and *B. pendula*, 18.08.2005, soil sampling.
- 5:** 51°19'28,5"N, 87°52'4,8"E, 853 m a.s.l., *A. sibirica* forest with *P. sibirica*, *Populus tremula* and *B. pendula*, 18.08.2005, soil sampling and hand sampling.
- 6:** 51°19'31,6"N, 87°52'16,1"E, 942 m a.s.l., *P. tremula* forest with *A. sibirica* and *P. sibirica*, 18.08.2005, soil sampling and hand sampling.
- 7:** 51°19'31,2"N, 87°52'21,1"E, 960 m a.s.l., *A. sibirica*, *P. sibirica* and *P. tremula* forest, 18.08.2005, soil sampling and hand sampling.
- 8:** 51°19'30,4"N, 87°52'50,0"E, 1100 m a.s.l., *P. sibirica* forest with *A. sibirica*, 17.08.2005, soil sampling.
- 8a:** 51°19'23,6"N, 87°53'2,1"E, 1191 m a.s.l., *P. sibirica* forest with *A. sibirica*, 17.08.2005, soil sampling and hand sampling.
- 9:** 51°19'07,5"N, 87°53'15,0"E, 1414 m a.s.l., *P. sibirica* forest with *A. sibirica*, 17.08.2005, soil sampling.
- 10:** 51°18'58,5"N, 87°53'33,3"E, 1468 m a.s.l., *P. sibirica* forest with *A. sibirica*, 17.08.2005, soil sampling.
- 10a:** 51°18'43,7"N, 87°54'23,7"E, 1699 m a.s.l., sparse growths of *P. sibirica* with *B. pendula* and *A. sibirica*, 16.08.2005, soil sampling.
- 11:** 51°18'41,3"N, 87°55'34,7"E, 1736 m a.s.l., old fire-site, *B. rotundifolia* and *Salix glauca* bushes with *P. sibirica* and *A. sibirica* sparse growths, 16.08.2005, soil sampling.
- 12:** 51°18'27,8"N, 87°54'57,4"E, 1847 m a.s.l., old fire-site, *B. rotundifolia* and *S. glauca* bushes with *P. sibirica* and *A. sibirica* sparse growths, 16.08.2005, soil sampling and hand sampling.
- 13:** 51°18'09,4"N, 87°55'43"E, 1861 m a.s.l., subalpine *P. sibirica* forest, 14.08.2005, hand sampling.
- 13a:** 51°18'24,1"N, 87°55'06,9"E, 1877 m a.s.l., subalpine sparse growths of *P. sibirica*, 16.08.2005, soil sampling and hand sampling.
- 14:** 51°18'23,0"N, 87°55'22,1"E, 1903 m a.s.l., subalpine sparse growths of *P. sibirica*, 14.08.2005, soil sampling.
- 15:** 51°18'24,5"N, 87°55'31,0"E, 1962 m a.s.l., subalpine sparse growths of *P. sibirica* with *A. sibirica*, 14.08.2005, soil sampling and hand sampling under stones.
- 16:** 51°18'33,6"N, 87°55'32,9"E, 2028 m a.s.l., subgoltsy sparse growths of *P. sibirica* with *B. rotundifolia* and *S. glauca* bushes, 15.08.2005, soil sampling and hand sampling.
- 18:** 51°18'30,0"N, 87°56'10,7"E, 2194 m a.s.l., *B. rotundifolia* and *S. glauca* mountain tundra with *Dryas* and lichens, 15.08.2005, soil sampling and hand sampling under stones.
- 19:** 51°18'30,5"N, 87°56'21,7"E, 2267 m a.s.l., summit of Malaya Koliusha Mt., *B. rotundifolia* and *S. glauca* rocky mountain tundra with *Dryas*, *Festuca* and lichens, 15.08.2005, soil sampling and hand sampling under stones.

Also we collected some material from two additional sampling sites (not included to official list of profile's sites):

- A:** 51°20'16,8"N, 87°51'47,6"E, about 500 m a.s.l., *Duscheikia fruticosa* forest on the bank of the river Bayas, 12.08.2005, hand sampling.
- B:** 51°18'05,4"N, 87°55'48,3"E, about 1900 m a.s.l., *P. sibirica* sparse growths, 14.08.2005, hand sampling.

The cluster analysis was performed using Statistica 10 (StatSoft 2011).

The material treated herein has been deposited mainly in the collection of the Altai State University, Barnaul, Russia (ASU), and partly shared also with the collection of the Institute of Biology and Soil Science, Far Eastern Branch, Russian Academy of Sciences, Vladivostok, Russia (IBSS), as indicated in the text. The species names include the literature references concerning Asian Russia only.

Taxonomic part

Order Julida Brandt, 1833

Family Julidae Leach, 1814

Genus *Julus* Linnaeus, 1758

Julus ghilarovi ghilarovi Gulička, 1963

Julus ghilarovi Gulička, 1963: 521, 520: figs.

Julus ghilarovi – Mikhaljova 2002: 206; Nefediev 2002b: 35; Mikhaljova and Nefediev 2003: 84.

Julus ghilarovi ghilarovi – Lokšina and Golovatch 1979: 386; Mikhaljova 1993: 11: figs; 2004: 59–61, 60: figs, 61: map; 2013a: 8; Mikhaljova and Golovatch 2001: 104; Nefediev 2005a: 41; 2005b: 8; Nefediev and Nefedieva 2006: 98; 2007a: 139; 2007b: 161; 2008a: 117; 2008b: 62; 2013: 87; Nefedieva and Nefediev 2008: 123; Babenko et al. 2009: 183; Nefediev et al. 2014: 63; Nefedieva et al. 2014: 65.

Material examined. 1 male, 7 females, 10 juv. (ASU), site 1; 1 male, 4 females, 3 juv. (ASU), site A; 1 female (IBSS), site 4; 5 males, 12 females, 4 juv. (ASU), site 5; 4 males, 6 females, 5 juv. (ASU), site 6; 1 female, 3 juv. (ASU), site 7; 3 juv. (ASU), site 8; 1 male, 1 female (IBSS), 3 males, 8 females, 11 juv. (ASU), site 19.

Distribution. This species appears to be widespread in the south of Siberia, Russia: Novosibirsk Area, Kemerovo Area, Altai Province, Republic of Altai, Republic of Khakassia, southern part of Krasnoyarsk Province. It is very likely it also occurs in the adjacent part of the Republic of Tyva.

Remarks. High ecological plasticity of this species allows it to inhabit different habitats like small-leaved, mixed and dark coniferous forests, herbaceous and alpine

meadows, and montane moss-stony tundras. In the Kyga Biogeocenosis Profile the animal prefers forest litter in low- and mid-mountain chern taiga forests up to about 1200 m a.s.l., and also recorded in rocky mountain tundra on the summit of Malaya Koliushka Mt. at 2267 m a.s.l. It is very likely that the species is displaced from subalpine sparse growths of *P. sibirica* by the congener of *Julus insolitus*. The numbers range from 0.5 to 18 ind./m².

Julus insolitus Mikhaljova, 2009

non *Julus gghilarovi brachydactylus* – Nefedieva and Nefediev 2008: 123.

Julus insolitus Mikhaljova, 2009b: 66–67, 64: figs.

Julus insolitus – Nefediev and Nefedieva 2013: 87; Nefedieva et al. 2014: 65.

Material examined. 4 males, 5 females, 4 juv. (ASU), site 9; 4 males, 1 female (IBSS), 6 males, 9 females, 10 juv. (ASU), site 10; 2 males, 6 females, 17 juv. (ASU), site 10a; 7 males, 17 females, 25 juv. (ASU), site 11; 20 males, 9 females, 49 juv., 1 fragm. (ASU), site 12; 1 male, 1 female, 1 juv. (ASU), site 13; 2 males, 4 females, 7 juv. (ASU), site 13a; 3 males, 8 females, 10 juv. (ASU), site 14; 1 female, 2 juv. (ASU), site 15; 14 males, 15 females, 22 juv. (ASU), site B; 10 males, 12 females, 15 juv. (ASU), site 16; 2 females, 6 juv. (ASU), site 18.

Distribution. The species appears to be spread only in the south of Siberia, Russia: Republic of Altai.

Remarks. This species has been described by Mikhaljova (2009b) on two male specimens, which are known to occur in forest-tundra and rocky tundra. In the Kyga Biogeocenosis Profile the species is collected from the upper line of mid-mountain chern taiga forests through subalpine sparse growths of *P. sibirica* to mountain tundra with *B. rotundifolia* and *S. glauca*. The maximum abundance is about 98 ind./m² registered in the old fire-site of sparse growths of *P. sibirica* now occupied with a succession of dwarf trees of *B. rotundifolia* and *S. glauca*. The above female specimens are the first records in this species.

Genus *Pacifiulus* Mikhaljova, 1982

Pacifiulus amurensis (Gerstfeldt, 1859)

Julus amurensis Gerstfeldt, 1859: 271.

Julus amurensis – Lokšina and Golovatch 1979: 387; Mikhaljova 1993: 34.

Pacifiulus imbricatus Mikhaljova, 1982: 211, 212: figs.

Pacifiulus imbricatus – Mikhaljova 1983: 87; 1988: 70; Mikhaljova and Petukhova 1983: 53; Ganin 1988: 7; 1989: 145; 1994: 60; 1995: 370; 1997: 10; Ryabinin et

al. 1988: 31; Mikhaljova and Bakurov 1989: 40; Gromyko 1990: 66; Mikhaljova 1993: 12; map; 1997: 145; Enghoff 1994: 27; Shelley et al. 2000: 50.

Pacifiulus amurensis – Mikhaljova 1998a: 5; 1998b: 64; figs, 65: map; 2004: 66–69, 67: figs, 68: map; 2009a: 603; 2009c: 3; 2012a: 23; 2012b: 112; Mikhaljova and Golovatch 2001: 105; Mikhaljova and Nefediev 2003: 84; Mikhaljova and Marusik 2004: 3; Nefediev 2005a: 48; 2005b: 8; Nefediev and Nefedieva 2006: 98; 2007a: 139; 2007b: 160; 2008b: 62; 2013: 87; Nefedieva et al. 2014: 65.

Material examined. 1 male (IBSS), site 9; 3 males (ASU), site 16; 1 male (ASU), site 18.

Distribution. This species is characterized by disjunctive area. The first distribution area is in the south of Siberia (Republic of Altai, Republic of Khakassia, southern part of Krasnoyarsk Province, Republic of Tyva) and the second one spreads in the Russian Far East (Maritime Province, southern part of Khabarovsk Province, Amur Area, Jewish Autonomous Area) and North-Eastern China.

Remarks. This species is characterized by euryoky, dwelling in Siberia in herbaceous meadows, small-leaved, mixed and dark coniferous forests, and subalpine meadows and montane tundras, up to 2500 m a.s.l. (Mikhaljova and Nefediev 2003). In the Kyga Biogeocenosis Profile the species is very rare (0.5–1.5 ind./m²), collected from the mid-mountain dark coniferous forest, and also from subgoltsy sparse growths of *P. sibirica* and mountain tundra with dwarf bushes of *B. rotundifolia* and *S. glauca*, with the maximum altitude registered is about 2194 m a.s.l.

Genus *Sibiriulus* Gulička, 1963

Sibiriulus altaicus Gulička, 1972

Cylindroiulus (Sibiriulus) altaicus Gulička 1972: 43–44, 44: fig.

Sibiriulus altaicus – Lokšina and Golovatch 1979: 387; Mikhaljova 1993: 13; 2004: 75: fig, 74: map; Mikhaljova and Golovatch 2001: 106; Mikhaljova et al. 2007: 57–59, 62, 58: figs; 2014: 45–47, 46: figs; Nefediev and Nefedieva 2007b: 162; 2008a: 117; 2008b: 62; 2013: 86–87; Nefedieva and Nefediev 2008: 123–124; Nefedieva et al. 2014: 65.

Material re-examined (specimens published by Mikhaljova et al. 2007, 2014). 1 male (ASU), site 1; 3 males, 2 juv. (IBSS), site 4; 1 male, 1 female, 2 juv. (IBSS), site 7; 1 male (ASU), site 8; 1 female (IBSS), 1 female (ASU), site 8A; 1 male (ASU), site 9; 1 male (IBSS), site 14.

Distribution. The species is known only in the south of Lake Teletskoye, Republic of Altai, its terra typica.

Remarks. In the Kyga Biogeocenosis Profile this species dwells in low- and mid-mountain chern taiga forests, and also subalpine sparse growths of *P. sibirica*, with the maximum altitude registered is about 1903 m a.s.l.

Family Nemasomatidae Bollman, 1893

Genus *Orinisobates* Lohmander, 1933

***Orinisobates sibiricus* (Gulička, 1963)**

Isobates sibiricus Gulička, 1963: 522: figs.

Isobates (Orinisobates) sibiricus – Gulička 1972: 45: figs; Nefediev and Nefedieva 2008a: 117.

Orinisobates sibiricus – Lokšina and Golovatch 1979: 387; Enghoff 1985: 53, 54: figs; Mikhaljova 1993: 16; 2002: 206; 2004: 96–97, 96: figs, 94: map; Mikhaljova and Golovatch 2001: 107; Mikhaljova and Nefediev 2003: 83; Nefediev 2005a: 39; 2005b: 8; Nefediev and Nefedieva 2006: 98; 2007a: 139; 2007b: 160; 2008a: 117; 2008b: 62; 2013: 87; Nefedieva and Nefediev 2008: 123; Nefediev et al. 2014: 63; Nefedieva et al. 2014: 65.

Material examined. 1 male (ASU), site 1; 1 female (ASU), site 8a.

Distribution. The species appears to be quite widespread in the south of Siberia, Russia: Kemerovo Area, Republic of Khakassia, Altai Province, Republic of Altai, southern part of Krasnoyarsk Province, Republic of Tyva, Chita Area. Also it has been recorded in Eastern Kazakhstan and Kyrgyzstan.

Remarks. The species inhabits forest litter of small-leaved, mixed and dark coniferous forests, under bark of logs and trees, and in mosses and mushrooms. The maximum altitude registered is about 1700 m a.s.l. (Mikhaljova and Golovatch 2001). In the Kyga Biogeocenosis Profile the species is very rare collected from low- and mid-mountain chern taiga forest, with the maximum abundance registered is about 1 ind./m².

Julidae gen. sp.

Material examined. 1 juv. (ASU), site 2; 2 females, 1 juv. (ASU), site 9; 1 female, 2 juv. (ASU), site 10.

Remarks. It is very likely these females and juveniles appear to belong to *S. altaicus* or *P. amurensis*.

Order Chordeumatida C. L. Koch, 1847

Family Diplomaragnidae Attems, 1907

Genus *Altajosoma* Gulička, 1972

***Altajosoma bakurovi bakurovi* (Shear, 1990)**

Diplomaragna bakurovi Shear, 1990: 22, 23: figs.

Diplomaragna bakurovi – Mikhaljova 1993: 18.

Altajosoma bakurovi – Mikhaljova 2000: 161; fig; 2004: 178–179, 178: figs, 116: map; Mikhaljova and Golovatch 2001:108; Nefediev 2002: 30; Mikhaljova and Nefediev 2003: 86; Mikhaljova et al. 2008: 51; Nefedieva and Nefediev 2008: 123–124; 2013: 87; Nefedieva et al. 2014: 65.

Material re-examined (specimen published by Mikhaljova et al. 2008). 1 male (ASU), site A.

Distribution. The species is known to occur in the south of Siberia, Russia: Tomsk, Novosibirsk and Kemerovo areas, Krasnoyarsk Province and Republic of Altai.

Remarks. This species dwells in various forest habitats like small-leaved, mixed and dark coniferous forests, and also mesophytous meadow and mountain tundra, with the maximum altitude registered is about 2500 m a.s.l (Mikhaljova and Nefediev 2003). In the Kyga Biogeocenosis Profile a single male is collected only by hand sampling from *D. fruticosa* forest on the bank of the river Bayas at about 500 m a.s.l.

Altajosoma deplanatum (Stuxberg, 1876)

Craspedosoma deplanatum Stuxberg, 1876a: 34, figs.

Craspedosoma deplanatum – Stuxberg 1876b: 317; Lokšina and Golovatch 1979: 382; Nefediev and Nefedieva 2008a: 117.

Altajosoma pinetorum Gulička, 1972: 37: figs.

Altajosoma pinetorum – Lokšina and Golovatch 1979: 382; Shelley et al. 2000: 62; Nefediev and Nefedieva 2008a: 117.

Diplomaragna deplanata – Shear 1990: 19, 20: figs; Mikhaljova 1993: 22.

Diplomaragna pinetorum – Shear 1990: 38; Mikhaljova 1993: 25.

Altajosoma deplanatum – Mikhaljova 2000: 160; 2004: 170–171, 171: figs, 162: map; 2013a: 7; Mikhaljova and Golovatch 2001: 108; Nefediev 2002b: 35; 2002d: 30; Mikhaljova and Nefediev 2003: 86; Nefediev 2005a: 50; 2005b: 9; Nefediev and Nefedieva 2005: 177; 2006: 98; 2007a: 139; 2007b: 161; 2007c: 99; 2008b: 62; 2011: 100; 2012a: 51; 2012b: 47; 2013: 87; Nefedieva and Nefediev 2008: 123; Mikhaljova et al. 2008: 51; Nefedieva et al. 2014: 65.

Material examined. 2 females (ASU), site 14.

Material re-examined (specimens published by Mikhaljova et al. 2008). 1 male (ASU), site 6; 1 male, 6 juv. (ASU), site 14.

Distribution. The species appears to be quite widespread in the south of Siberia, Russia: Tomsk, Novosibirsk and Kemerovo areas, Republic of Altai and Republic of Khakassia, and originally described from between the city of Achinsk (Krasnoyarsk Province) and the city of Mariinsk (Kemerovo Area).

Remarks. This species lives mainly in various forest habitats like small-leaved, mixed and dark coniferous forests, forest- and shrub tundra, and also mesophytous meadow. The maximum altitude registered is about 2080 m a.s.l (Mikhaljova et al.

2008). In the Kyga Biogeocenosis Profile the species quite rare found in the mid-mountain mixed forest and in the subalpine sparse growths of *P. sibirica*, with the maximum abundance registered is about 6 ind./m².

Altajosoma katunicum Mikhaljova, 2000

Altajosoma katunicum Mikhaljova, 2000: 161–162, 162: figs.

Altajosoma katunicum – Mikhaljova and Golovatch 2001: 108; Mikhaljova 2004: 176–177, 177: figs, 112: map; Mikhaljova et al. 2008: 52; Nefedieva et al. 2014: 65.

Material re-examined (specimens published by Mikhaljova et al. 2008). 1 male (ASU), site 1.

Distribution. This species is known to occur only in the Republic of Altai, Russia.

Remarks. This species originally described from the Katunskii Mt. Range, Central Altai at 1600–2200 m a.s.l. (Mikhaljova 2000). In the Kyga Biogeocenosis Profile the species is very rare (1 ind./m²), collected only from the mixed forest in the lowest study site at 443 m a.s.l.

Altajosoma kemerovo (Shear, 1990)

Diplomaragna kemerovo Shear, 1990: 21, 20: figs.

Diplomaragna kemerovo – Mikhaljova 1993: 25; Nefediev and Nefedieva 2008a: 117.

Altajosoma kemerovo – Mikhaljova 2000: 161; 2004: 180–181, 180: figs, 173: map; 2013a: 7; Mikhaljova and Golovatch 2001: 108; Vorobiova et al. 2002: 60; Mikhaljova and Nefediev 2003: 86; Nefediev 2005a: 53; 2005b: 9; Nefediev and Nefedieva 2006: 98; 2007a: 139; 2007b: 161; 2008b: 62; 2013: 87; Nefedieva and Nefediev 2008: 123; Mikhaljova et al. 2008: 51; Nefedieva et al. 2014: 65.

Material examined. 4 females (ASU), site 7; 1 juv. (ASU), site 8; 7 juv. (ASU), site 13a; 1 female, 3 juv. (ASU), site 16.

Material re-examined (specimens published by Mikhaljova et al. 2008). 1 male (ASU), site A; 1 male (ASU), site 5; 1 male (ASU), site 6; 5 males (ASU), site 7; 1 male (ASU), site 8; 1 male (ASU), site 13a; 1 male (ASU), site 16.

Distribution. This species appears to be spread in the south of Siberia, Russia: Kemerovo and Novosibirsk areas, Republic of Altai, Republic of Khakassia, southern part of Krasnoyarsk Province.

Remarks. The species dwells in different forest habitats like small-leaved, mixed and dark coniferous forests. In the Kyga Biogeocenosis Profile it is very rare (0.5–5 ind./m²), mainly collected in low- and mid-mountain chern taiga at 853–1100 m

a.s.l., but also found in subalpine sparse growths of *P. sibirica* and subgoltsy sparse growths of *P. sibirica* with *B. rotundifolia* and *S. glauca* bushes, with the maximum altitude registered is about 2028 m a.s.l.

***Shearia teletskaya* Mikhaljova, 2000**

Shearia teletskaya Mikhaljova, 2000: 167–168, 167: figs.

Shearia teletskaya – Mikhaljova and Golovatch 2001: 111; Mikhaljova 2004: 160–161, 160: figs, 112: map; Mikhaljova et al. 2008: 54; Nefedieva et al. 2014: 65.

Material examined. 1 juv. (ASU), site 16.

Material re-examined (specimens published by Mikhaljova et al. 2008). 1 male, 4 juv. (ASU), site B; 1 male, 8 juv. (ASU), site 12; 1 male (IBSS), 1 male (ASU), site 16.

Distribution. The species is known to occur only in the south of Lake Teletskoye, Republic of Altai, Russia.

Remarks. This species inhabits dark coniferous taiga forests at 1350–1750 m a.s.l. and the subalpine belt (= goltsy) at 1750–2000 m a.s.l. (Mikhaljova 2004). In the Kyga Biogeocenosis Profile sites investigated the species is very rare (3 ind./m²), mainly collected by hand sampling in subalpine sparse growths of *P. sibirica* at 1847–2028 m a.s.l.

Diplomaragnidae gen. sp.

Material examined. 1 female (ASU), site A; 1 female, 1 juv. (ASU), site 5; 5 juv. (ASU), site 8a; 4 juv. (ASU), site 9; 1 female, 14 juv. (ASU), site 10; 5 juv. (ASU), site 10a; 1 female, 9 juv. (ASU), site 11; 1 female, 1 fragm. (ASU), site 13; 1 female, 1 juv. (ASU), site 18; 1 female, 1 juv. (ASU), site 19.

Remarks. The above specimens appear to belong to some species of *Altajosoma* or *S. teletskaya*.

Family Anthroleucosomatidae Verhoeff, 1899

Genus *Ghilarovia* Gulička, 1972

***Ghilarovia kygae* Gulička, 1972**

Ghilarovia kygae Gulička, 1972: 39, 40: figs.

Ghilarovia kygae – Lokšina and Golovatch 1979: 383; Shear 1988: 55: figs; Mikhaljova 1993: 16; 2002: 203, 202: figs; 2004: 188–190, 188: figs, 189: map; 2013a: 8; Shelley et al. 2000: 68; Mikhaljova and Golovatch 2001: 107; Nefediev 2005a: 58; 2005b: 9; Nefediev and Nefedieva 2007b: 161; 2008a: 117; 2008b: 62; 2013: 87; Nefedieva and Nefediev 2008: 123; Nefedieva et al. 2014: 65.

Material examined. 18 males, 16 females (ASU), site 1; 9 males, 6 females (ASU), site 2; 2 males, 1 female (ASU), site A; 3 males, 9 females, 2 juv. (ASU), site 4; 24 males, 18 females, 2 juv. (ASU), site 5; 15 males, 15 females, 2 juv. (ASU), site 6; 21 males, 15 females, 1 juv., 1 fragm. (ASU), site 7; 5 males, 3 females (ASU), site 8; 6 males, 6 females, 14 juv. (ASU), site 8a; 2 females, 1 juv. (ASU), site 9; 2 males (ASU), site 10a; 3 males, 3 females (ASU), site 11; 3 males, 1 female, 8 juv. (ASU), site 12; 3 males, 1 female (ASU), site 12; 6 males, 8 females (ASU), site 13; 2 males, 5 females (ASU), site 13a; 5 males, 16 females (ASU), site 14; 1 male, 1 female, 1 fragm. (ASU), site 15; 2 males, 8 females (ASU), site B; 1 male (ASU), site 16.

Distribution. The species is known to occur in the Republic of Altai, Siberia, Russia only.

Remarks. This species prefers to live in mixed and dark coniferous forests, and also known from subalpine habitats like golsty, with the maximum altitude registered is about 2000 m a.s.l. (Mikhaljova and Golovatch 2001). In the Kyga Biogeocenosis Profile the species shows the highest ecological plasticity, dwelling in low- and mid-mountain chern taiga forests, and subalpine sparse growths of *P. sibirica* up to subgolsty, with the maximum altitude registered is about 2028 m a.s.l.

Family Kirkayakidae Özdikmen, 2008

(syn. Altajellidae Mikhaljova & Golovatch, 2001)

Genus *Kirkayakus* Özdikmen, 2008

(syn. *Altajella* Gulička, 1972)

***Kirkayakus pallidus* (Gulička, 1972)**

(syn. *Altajella pallida* Gulička, 1972)

Altajella pallida Gulička, 1972: 42, 43: figs.

Altajella pallida – Lokšina and Golovatch 1979: 383; Shear 1988: 51; Mikhaljova 1993: 34; 2004: 196–199, 197: figs, 105: map; Shelley et al. 2000: 61; Mikhaljova and Golovatch 2001: 111, 112: figs; Nefediev 2005a: 58; 2005b: 9; Nefediev and Nefedieva 2007b: 161; 2008a: 117; 2008b: 62; Nefedieva and Nefediev 2008: 123.

Kirkayakus pallidus – Özdikmen 2008: 342; Nefediev and Nefedieva 2013: 87; Nefedieva et al. 2014: 65.

Material examined. 1 male (ASU), site 8; 1 female (ASU), site 9.

Distribution. This species is an endemic in the south of Lake Teletskoye, Republic of Altai, Siberia, Russia.

Remarks. The species lives in dark coniferous forests up to 1350 m a.s.l. (Mikhaljova and Golovatch 2001). In the Kyga Biogeocenosis Profile the animal prefers mid-mountain dark coniferous forests at 1100–1414 m a.s.l., when it is very rare, and the numbers range from 0.5 to 1 ind./m². The above material appears to belong to topotypes, and this is the first record of the female specimen in this species.

Genus *Teleckophoron* Gulička, 1972

Teleckophoron montanum Gulička, 1972

Teleckophoron montanum Gulička, 1972: 41: figs.

Teleckophoron montanum – Lokšina and Golovatch 1979: 383; Mikhaljova 1993: 35; 2004: 193–196, 195: figs, 107: map; Shelley et al. 2000: 79; Mikhaljova and Golovatch 2001: 113, 114: figs; Nefediev 2005a: 59; 2005b: 9; Nefediev and Nefedieva 2006: 98; 2007b: 161; 2008a: 117; 2008b: 62; 2013: 87; Nefedieva and Nefediev 2008: 123; Nefedieva et al. 2014: 65.

Material examined. 3 males, 1 female, 4 juv. (ASU), site 8; 1 female, 3 juv. (ASU), site 8a.

Distribution. The area of this species appears to encompass the Republic of Altai and the southern part of the Krasnoyarsk Province, both Siberia, Russia.

Remarks. This species inhabits dark coniferous forests and montane tundras. The maximum altitude registered is about 1000 m a.s.l. (Gulička 1972). In the Kyga Biogeocenosis Profile the species prefers mid-mountain dark coniferous forests up to 1191 m a.s.l., where the numbers range from 3 to 8 ind./m².

Order Polydesmida Leach, 1815

Family Polydesmidae Leach, 1815

Genus *Schizoturanius* Verhoeff, 1931

Schizoturanius clavatipes (Stuxberg, 1876)

Polydesmus clavatipes Stuxberg, 1876a: 34, figs.

Polydesmus clavatipes – Stuxberg 1876b: 316; Nefediev and Nefedieva 2008a: 117.

Schizoturanius clavatipes – Lohmander 1933: 27; Hoffman 1975: 81, 82: figs; Lokšina and Golovatch 1979: 384; Mikhaljova 1993: 31, 32: figs; 2002: 206; 2004: 238–240, 239: figs, 228: map; 2013a: 9; 2013b: 221; Nefediev 2001: 84; 2002c: 139; 2002d: 30; Mikhaljova and Golovatch 2001: 116; Vorobiova et al. 2002: 60; Mikhaljova and Nefediev 2003: 81; Nefediev 2005a: 61; 2005b: 9; Nefediev and Nefedieva 2005: 178; 2006: 98; 2007a: 139; 2007b: 161; 2007c: 102; 2008b: 62; 2011: 100; 2012a: 51; 2012b: 47; 2013: 87;

Nefedieva and Nefediev 2008: 123; Nefediev et al. 2014: 63; Nefedieva et al. 2014: 65.

Material examined. 4 males, 2 females, 4 juv. (ASU), site 1; 2 males, 2 females, 3 juv. (ASU), site 2; 2 males, 1 female, 1 juv. (ASU), site 4; 4 males, 2 females, 1 juv. (ASU), site 5; 2 males, 1 females (ASU), site 6; 6 males, 2 females, 11 juv. (ASU), site 7; 1 male, 1 female (ASU), site 8a; 1 female (ASU), site 9; 7 juv. (ASU), site 14; 3 juv. (ASU), site B.

Distribution. This species appears to be quite widespread in the south of Siberia, Russia: Tomsk, Novosibirsk and Kemerovo areas, Altai Province, Republic of Altai, Republic of Khakassia, southern part of Krasnoyarsk Province.

Remarks. Being highly euryoecic, the species populates various forest habitats (small-leaved, mixed and dark coniferous forests), and also meadows and glades. In the Kyga Biogeocenosis Profile the species prefers low- and mid-mountain chern taiga forests, where its numbers range from 0.5 to 10 ind./m², but also collected from subalpine sparse growths of *P. sibirica* at about 1903 m a.s.l.

Schizoturanius tabescens (Stuxberg, 1876)

Polydesmus tabescens Stuxberg, 1876a: 35, figs.

Polydesmus tabescens – Stuxberg 1876b: 316; Lokšina and Golovatch 1979: 385.

Turanodesmus salairicus Gulička, 1963: 523, 522: figs; Nefediev and Nefedieva 2008a: 117.

Schizoturanius salairicus – Lokšina and Golovatch 1979: 384; Mikhaljova 1993: 31; Nefediev 2001: 84, 2002a: 40; 2002b: 35; 2002c: 139; 2002d: 30; Mikhaljova and Golovatch 2001: 116; Mikhaljova and Nefediev 2003: 83.

Schizoturanius tabescens – Mikhaljova 1993: 31, 32: figs; 2004: 240–242, 241: figs, 242: map; 2013b: 221; Vorobiova 1999: 33; Mikhaljova and Golovatch 2001: 116; Vorobiova et al. 2002: 60; Rybalov 2002; Mikhaljova and Marusik 2004: 8, 7: figs; Nefediev 2005a: 64; 2005b: 9; Nefediev and Nefedieva 2005: 178; 2006: 98; 2007a: 139; 2007b: 161; 2007c: 102; 2008a: 117; 2008b: 62; 2011: 100; 2012a: 51; 2012b: 47; 2013: 87; Nefedieva and Nefediev 2008: 123; Babenko et al. 2009: 183; Nefedieva et al. 2014: 65.

Material examined. 3 juv. (ASU), site 2; 2 females, 2 juv., 1 fragm. (ASU), site 4; 4 juv. (ASU), site 5; 17 juv. (ASU), site 7; 2 females, 1 juv. (ASU), site 8a; 1 female (ASU), site 9; 2 females (ASU), site B.

Distribution. Being rather widespread, the species is known to occur in the south of Siberia, Russia: Tomsk, Novosibirsk and Kemerovo areas, Altai Province, Republic of Altai, Republic of Khakassia, southern part of Krasnoyarsk Province.

Remarks. This species lives in different forests like small-leaved, mixed and dark coniferous ones, and also populates meadows. In the Kyga Biogeocenosis Profile the animal prefers low- and mid-mountain chern taiga forests, where its numbers range from 0.5 to 8 ind./m², but also collected from subalpine sparse growths of *P. sibirica* at about 1900 m a.s.l.

Results

The millipede diversity in the south of Lake Teletskoye is estimated to be at least 15 species and subspecies from 10 genera, 6 families and three orders: *Julus ghilarovi ghilarovi* Gulička, 1963, *J. insolitus* Mikhaljova, 2009, *Orinisobates sibiricus* (Gulička, 1963), *Pacifiulus amurensis* (Gertsfeldt, 1859), *Sibiriulus altaicus* (Gulička, 1972), *Ghilarovia kygae* Gulička, 1972, *Kirkayakus pallidus* (Gulička, 1972) (synonym of *Altajella pallida* Gulička, 1972), *Teleckophoron montanum* Gulička, 1972, *Altajosoma bakurovi bakurovi* (Shear, 1990), *A. deplanatum* (Stuxberg, 1876), *A. katunicum* Mikhaljova, 2000, *A. kemerovo* (Shear, 1990), *Shearia teletskaya* Mikhaljova, 2000, *Schizoturanius clavatipes* (Stuxberg, 1876) and *S. tabescens* (Stuxberg, 1876).

The bulk of species diversity is confined both to low- and mid-mountain chern taiga forests and high-mountain shrub tundras with *Betula rotundifolia* and *Salix glauca*, achieving from 5 to 9 species, whereas subalpine sparse growths of *Pinus sibirica* are characterized by the lowest millipede diversity, with 3 the most widespread species (Table 1). One of them, *G. kygae*, has the maximum of ecological plasticity. This species is recorded in almost all study sites, dwelling in low- and mid-mountain chern taiga forests, and subalpine sparse growths of *P. sibirica* up to subgoltsy, with the maximum altitude registered is about 2028 m a.s.l.

The numbers of diplopods range from 14.5 to 65 ind./m² in subzones of low- and mid-mountain chern taiga forests, and from 5.5 to 8 ind./m² in high-mountain shrub tundras (Table 1). Despite of the lowest species diversity in subalpine sparse growths of *P. sibirica*, millipedes show the maximum numbers, reaching up to 130 ind./m² in the old fire-site (site 11), evidently caused by the abundance of plant debris of dwarf vegetation of *Betula rotundifolia* and *Salix glauca* that appear to be more suitable for feeding of millipedes than pine litter.

Clustering of investigated sites in the Kyga Biogeocenosis Profile on species diversity allows to grouping at least two obvious sets of sites (Figure 1). The first group unites low-mountain chern taiga forests (sites from 2 to 7), with altitudes range from 494 to 960 m a.s.l. The second group includes subalpine sparse growths of *P. sibirica* localities (sites from 10a to 15), and also the highest locality of chern taiga forest (site 10) and a transition locality from subalpine sparse growths to shrub mountain tundra (site 16), with altitudes range from 1468 to 2028 m a.s.l. Some localities of mid-mountain chern taiga forests and mountain tundras are less similar both to each other and to other groups.

The very interesting situation is observed with two congeners of *Julus* as regards to altitude distribution, showing competitive relationships to each other. Thus, *J. g. ghilarovi* is mainly recorded in low-mountain and in the beginning of mid-mountain chern taiga forests, while it almost disappears in subalpine sparse growths of *P. sibirica*, but it emerges again in rocky mountain tundra on the top of the investigated biogeocenosis profile. At the same time, *J. insolitus* populates mainly subalpine sparse growths of *P. sibirica* and lifts up to shrub mountain tundra at 2194 m a.s.l. Taking into account that *J. insolitus* is very abundant here, we assume it wins the competition and

Table 1. Millipede abundance (ind./m²) and hand sampling (+) in the Kyga Biogeocoenosis Profile sites investigated.

Species	Sites																							
	1	2	A	4	5	6	7	8	8a	9	10	10a	11	12	13	13a	B	14	15	16	18	19		
Julida																								
<i>J. g. ghilarovi</i>	18		+	0.5	3	5	2	3															6	
<i>J. insolitus</i>										6.5	15	50	98	20	+	13	+	14	3	15.5	4			
<i>S. altaicus</i>	1			2.5			2	1	2	0.5								1						
<i>O. sibiricus</i>	1								1															
<i>P. amurensis</i>										0.5											1.5	0.5		
Julidae gen. sp.		1								1.5	1.5													
Chordeumatida																								
<i>A. b. bakurovi</i>			+																					
<i>A. deplanatum</i>						+												6						
<i>A. katunicum</i>	1																							
<i>A. kemeroovo</i>			+		1.5	0.5	3	2								5				2.5				
<i>S. teletskaya</i>														3			+			+				
Diplomaragnidae gen. sp.			+						4	2	7.5	10	20		+							1	2	
<i>G. kygae</i>	34	15	+	7	16.5	14.5	14.5	8	22	1.5	4	12	2	+	4	+	14	2	0.5					
<i>K. pallidus</i>								1		0.5														
<i>T. montanum</i>								8	3															
Polydesmida																								
<i>S. clavipes</i>	10	7		2	1.5	1	8		2	0.5							+	4.5						
<i>S. tabescens</i>		3		2.5	2		8		2	0.5							+							
Total abundance	65	26	-	14.5	24.5	21	37.5	23	36	14	24	64	130	25	-	22	-	39.5	5	20	5.5	8	8	
Numbers of species	6	4	4	4	5	5	5	6	6	7	9	3	3	3	3	3	5	5	2	5	2	5	3	2




Figure 1. Neighbour-joining tree of similarity / dissimilarity of studied sites on species diversity.

displaces *J. g. ghilarovi* from subalpine *P. sibirica* sparse growths. The record of female specimens of *J. insolitus* is the first one in this species.

Both members of Kirkayakidae, *K. pallidus* and *T. montanum*, are reported in chern mid-mountain taiga at the highest altitudes for the first time. Also this is the first record of the female specimen in the former species.

Acknowledgements

We are thankful to D.A. Demidko (Krasnoyarsk, Russia) and V.S. Bogdanov (Chiri, Russia) for their great help with the organization of the collecting trip in Altai for the first and second authors. Special thanks are also addressed to anonymous reviewers and section editor for comments.

References

- Babenko AS, Nefediev PS, Nefedieva JS (2009) [The fauna and population dynamics of millipedes (Diplopoda) of chern taiga of Salair]. Vestnik Tomskogo gosudarstvennogo universiteta, Seriya biologiya 319: 182–185. [In Russian]
- Enghoff H (1985) The millipede family Nemasomatidae with the description of a new genus and a revision of *Orinisobates* (Diplopoda, Julida). Entomologica Scandinavica 16: 27–67. doi: 10.1163/187631285X00045

- Enghoff H (1994) Geographical parthenogenesis in millipedes (Diplopoda). *Biogeographica* 70(1): 25–31.
- Ganin GN (1988) [Estimation of the role of millipedes of Siberian pine-broadleaved forests of Priamurye in litter destruction]. *Izvestiya Sibirskogo otdeleniya AN SSSR, Biology* 20(3): 7–10. [In Russian, with English summary]
- Ganin GN (1989) [The role of millipedes in the decay and transformation of forest litter in the southern Khabarovsk Province]. *Zoologicheskii Zhurnal* 68(1): 145–149. [In Russian, with English summary]
- Ganin GN (1994) [Role of soil mesofauna in the biogenic turnover of elements in forest and meadow ecosystems of the Amur region]. *Ekologiya* 5–6: 59–67. [In Russian]
- Ganin GN (1995) [Heavy metals in soil invertebrates in nature reserves of the Russian Far East]. *Ekologiya* 5: 368–372. [In Russian]
- Ganin GN (1997) [Soil animals of the Ussuri region]. Dalnauka Publ., Vladivostok-Khabarovsk, 159 pp. [In Russian, with English summary]
- Gerstfeldt G (1859) Ueber einige zum Theil neue Arten Platoden, Anneliden, Myriapoden und Crustaceen Sibiriens, namentlich seines östlichen Theiles und des Amur-Gebietes. *Mém. savants étrangers Acad. Imp. Sci. St. Pétersbourg* 8: 1–36.
- Ghilarov MS (1987) [Recording of large invertebrates (mesofauna)]. In: *Quantitative methods in soil zoology*. Nauka Publ., Moscow, 9–26. [In Russian]
- Gromyko MN (1990) [Peculiarities in the structure of soil populations in oakwood ecosystems of the Sikhote-Alin Biosphere Reserve]. In: *Ecological investigations in the Sikhote-Alin Nature Reserve*. TSNIL Glavokhoty RSFSR Publ., Moscow, 57–72. [In Russian]
- Gulička J (1963) [New millipedes (Diplopoda) from the USSR. Part 1]. *Zoologicheskii Zhurnal* 42(4): 518–524. [In Russian, with English summary]
- Gulička J (1972) [New millipedes (Diplopoda) from the USSR. Part 2]. *Zoologicheskii Zhurnal* 51(1): 36–45. [In Russian, with English summary]
- Hoffman RL (1975) A note on the status of *Polydesmus clavatipes* and some related species from Central Asia (Diplopoda, Polydesmidae). *Zoologischer Anzeiger* 194: 79–83.
- Lohmander H (1933) Über Diplopoden aus Zentralasien. *Arkiv för Zoologi* 25A(6): 1–71. [for 1932]
- Lokšina IE, Golovatch SI (1979) Diplopoda of the USSR fauna. *Pedobiologia* 19(6): 381–389.
- Mikhailjova EV (1982) [New millipedes of the family Julidae (Diplopoda) from the USSR Far East]. *Zoologicheskii Zhurnal* 61(2): 210–216. [In Russian, with English summary]
- Mikhailjova EV (1983) [Distribution of millipedes (Diplopoda) along the soil profile in the forests of the Primorsky Province]. In: *Fauna and ecology of arthropods in the Far East*. Vladivostok, 77–90. [In Russian]
- Mikhailjova EV (1988) [Millipedes (Diplopoda) of the Ussuriyskii and «Kedrovaya Pad» nature reserves]. In: *A role of insects in biocenoses of the Far East*. Vladivostok, 114–126. [In Russian]
- Mikhailjova EV (1993) The millipedes (Diplopoda) of Siberia and the Far East of Russia. *Arthropoda Selecta* 2(2): 3–36.
- Mikhailjova EV (1998a) On new and poorly-known millipedes (Diplopoda) from the Far East of Russia. *Far Eastern Entomologist* 60: 1–8.

- Mikhailjova EV (1998b) The millipedes of the Far East of Russia (Diplopoda). *Arthropoda Selecta* 7(1): 1–77.
- Mikhailjova EV (2000) Review of the millipede family Diplomaragnidae (Diplopoda: Chordeumatida). *Arthropoda Selecta* 8(3): 153–181. [for 1999]
- Mikhailjova EV (2002) On some poorly-known millipedes from Siberia (Diplopoda). *Arthropoda Selecta* 10(3): 201–207. [for 2001]
- Mikhailjova EV (2004) The millipedes (Diplopoda) of the Asian part of Russia. Pensoft Publishers, Sofia-Moscow, Series Faunistica 39, 292 pp. doi: 10.11646/zootaxa.3866.1.2
- Mikhailjova EV (2009a) The millipedes (Diplopoda) of the Russian Far East islands and the Kamchatka Peninsula. *Soil Organisms* 81(3): 599–616.
- Mikhailjova EV (2009b) New species of the family Julidae Leach 1814 from Altai, Russia (Diplopoda, Julida). *Zootaxa* 2235: 59–68.
- Mikhailjova EV (2009c) A list of the millipedes (Diplopoda) of the Russian Far East. *Far Eastern Entomologist* 197: 1–8.
- Mikhailjova EV (2012a) [Class Diplopoda – Millipedes]. *Wildlife of Bastak Nature Reserve. Blagoveshchensk State Pedagogical University Publ., Blagoveshchensk*, 22–25. [In Russian]
- Mikhailjova EV (2012b) [Millipede (Diplopoda) taxonomic diversity in the Asian part of Russia]. In: *Regions of a new exploration: theoretical and practical questions of biodiversity and landscape investigations and conservation. Institute for Water and Environmental Problems Publ., Khabarovsk*, 111–114. [In Russian, abstract in English]
- Mikhailjova EV (2013a) New data on the millipede fauna (Diplopoda) of Altai, Russia. *Far Eastern Entomologist* 265: 1–10.
- Mikhailjova EV (2013b) [Endemism of millipedes (Diplopoda) in the Asian part of Russia]. Investigation, conservation and restoration of natural landscapes. In: *Proceedings of III International Scientific Conference, Volgograd. Planeta Publ., Moscow*, 220–223. [In Russian]
- Mikhailjova EV, Bakurov VD (1989) [On millipedes (Diplopoda) of the Siniy Mountain Ridge (Sikhote-Alin)]. In: *Soil invertebrates in the south of the Far East. Khabarovsk*, 38–47. [In Russian]
- Mikhailjova EV, Golovatch SI (2001) A review of the millipede fauna of Siberia (Diplopoda). *Arthropoda Selecta* 9(2): 103–118. [for 2000]
- Mikhailjova EV, Marusik YM (2004) New data on taxonomy and fauna of the millipedes from the Russian Far East, Siberia and Mongolia (Diplopoda). *Far Eastern Entomologist* 133: 1–12.
- Mikhailjova EV, Nefediev PS (2003) A contribution to the millipede fauna of Siberia (Diplopoda). *Arthropoda Selecta* 11(1): 81–87. [for 2002]
- Mikhailjova EV, Nefediev PS, Nefedieva JS (2007) New data on millipedes of the family Julidae (Diplopoda, Julida) from Altai, Siberia. *Zootaxa* 1541: 57–63.
- Mikhailjova EV, Nefediev PS, Nefedieva JS (2008) A new species and new records of millipedes of the family Diplomaragnidae (Diplopoda, Chordeumatida) from Altai. *Zootaxa* 1931: 49–56.
- Mikhailjova EV, Nefediev PS, Nefedieva JS, Sakhnevich MB, Dyachkov YuV (2014) Review of the millipede genus *Sibiriulus* Gulička, 1972, with descriptions of three new species from Altai, Siberia, Russia (Diplopoda, Julida, Julidae). *Zootaxa* 3866(1): 30–52.

- Mikhailjova EV, Petukhova EL (1983) [A comparative analysis of the millipede faunas (Diplopoda) of Primorsky Province's forests with the help of inclusion and similarity indices]. In: Graph theoretic methods in biogeographical investigations. Vladivostok, 48–66. [In Russian]
- Nefediev PS (2001) [On the fauna and ecology of myriapods (Myriapoda) in the environs of the village of Smolenskoe, Altai Province]. In: Landscapes of Western Siberia: Investigation problems, ecology and rational use. Proceedings of 7 International Conference, devoting to the International Day of the Earth. Biysk Pedagogical State University Publ., Biysk, 84–86. [In Russian]
- Nefediev PS (2002a) [Eco-faunistic investigations of myriapods in the Teguldet District, Tomsk Area]. In: Proceedings of International Conference of Students and Young Researchers «Lomonosov–2002». Moscow State University Publ., Moscow, 7: 40–41. [In Russian]
- Nefediev PS (2002b) [The fauna and ecology of myriapods (Myriapoda) of a relict lime grove (village of Kuzedeyevo)]. In: Proceedings of 40 International Scientific Student Conference «Students and Scientific and Technical Progress». Novosibirsk State University Publ., Novosibirsk, 35. [In Russian]
- Nefediev PS (2002c) [Populations and some ecological peculiarities of myriapods of gray forest soils in the southern of the Tomsk Area]. In: Biology – Science of the XXI Century. Abstracts of 6 School Conference of Young Researchers. Pushchino Scientific Centre RAS Publ., Pushchino-on-Oka, 138–139. [In Russian]
- Nefediev PS (2002d) On the Diplopoda fauna of South-West Siberia. In: 12th International Congress of Myriapodology. Book of Abstracts. Pietermaritzburg, 30.
- Nefediev PS (2005a) [Millipedes (Myriapoda, Diplopoda) of the Southeast of Western Siberia (fauna, zoogeography, ecology)]. Thesis of Candidate (Ph.D.) of Biological Sci. Degree. Tomsk, 175 pp. [In Russian]
- Nefediev PS (2005b) [Millipedes (Myriapoda, Diplopoda) of the Southeast of Western Siberia (fauna, zoogeography, ecology)]. Autoreferate of the Thesis of Candidate (Ph.D.) of Biological Sci. Degree. UFO-print Publ., Tomsk, 21 pp. [In Russian]
- Nefediev PS, Dyachkov YuV, Nefedieva JS (2014) Fauna and ecology of millipedes (Diplopoda) in the Tigirek State Nature Reserve, Russian Altai. In: Tuf IH, Tajovský K (Eds) 16th International Congress of Myriapodology. Book of Abstracts. Institute of Soil Biology, BC ASCR & Faculty of Science, Palacký University, Olomouc, 63.
- Nefediev PS, Nefedieva JS (2005) [Seasonal fluctuations of diplopod density in some forests of Western Siberia]. In: Ecological diversity of soil biota and biological productivity of soils. Proceedings of IV (XIV) All-Russian Conference of Soil Zoology. Tyumen State University Publ., Tyumen, 177–178. [In Russian]
- Nefediev PS, Nefedieva JS (2006) [Regional peculiarities of millipede fauna (Diplopoda) in the south-east of Western Siberia]. In: Ecology of South Siberia and adjacent territories. Proceedings of X International School Conference of Students and Young Researchers. Khakassian State University Publ., Abakan 1: 98. [In Russian]
- Nefediev PS, Nefedieva JS (2007a) A brief analysis of the biotopic distribution of millipedes (Diplopoda) in the south-east of Western Siberia. In: Forest Soils: Research Results, Problems and Future Outlook. Abstracts of International Conference. Institute of Biology, Komi Scientific Center, Ural Division of RAS Publ., Syktyvkar, 139–140. [In Russian and in English]

- Nefediev PS, Nefedieva JS (2007b) [Biogeographical characteristic of millipede fauna in the south-east of Western Siberia]. In: Biodiversity of invertebrate animals. Proceedings of II All-Russian School Conference «Conceptual and applied aspects of scientific investigations and education in invertebrate zoology». Deltaplan Publ., Tomsk, 133–138. [In Russian]
- Nefediev PS, Nefedieva JS (2007c) [Seasonal dynamics of locomotor activity of millipedes (Diplopoda) in forests of Western Siberia]. In: Ecological problems of unique natural and anthropogenic landscapes. Proceedings of All-Russian Conference. Yaroslavl State University, Yaroslavl, 98–103. [In Russian]
- Nefediev PS, Nefedieva JS (2008a) [A historical review of faunistic investigations of millipedes (Diplopoda) in Western Siberia]. In: Altai: ecology and environmental management. Proceedings of VII Russian-Mongolian Conference of Students and Young Researchers. Biysk Pedagogical State University Publ., Biysk 1: 117–120. [In Russian]
- Nefediev PS, Nefedieva JS (2008b) Zoogeographical analysis of the millipede fauna (Diplopoda) in the south-east of Western Siberia. In: Myriapoda and Onychophora of the World Diversity, Biology and Importance. Abstracts of 14th International Congress of Myriapodology. Staatliches Museum für Naturkunde, Görlitz. Peckiana 6: 62.
- Nefediev PS, Nefedieva JS (2011) [Millipedes (Diplopoda) of green plantations of Tomsk City and its suburbs]. In: Conceptual and applied aspects of scientific investigations in invertebrate zoology. Proceedings of III All-Russian School Conference. Agraf-Press Publ., Tomsk, 100–102. [In Russian]
- Nefediev PS, Nefedieva JS (2012a) [Some peculiarities of distribution of millipedes (Diplopoda) along the soil profile in the south-taiga and small-leaved forests of Western Siberia]. *Izvestia Altaiskogo gosudarstvennogo universiteta, Biologicheskie nauki* 3(75),1: 49–54. [In Russian]
- Nefediev PS, Nefedieva JS (2012b) [Some peculiarities of seasonal dynamics of sex-age structure of milliped populations (Diplopoda) in the southern taiga and small-leaved forests of Western Siberia]. *Izvestia Altaiskogo gosudarstvennogo universiteta, Biologicheskie nauki* 3(75),2: 46–48. [In Russian]
- Nefediev PS, Nefedieva JS (2013) [Biodiversity and ecology of millipedes in the environs of Lake Teletskoe (Diplopoda)]. *Izvestia Altaiskogo gosudarstvennogo universiteta, Biologicheskie nauki* 3(79),1: 86–87. [In Russian]
- Nefedieva JS, Nefediev PS (2008) Ecofaunistic investigations of millipedes (Diplopoda) in the environs of Lake Teletskoe. In: Myriapoda and Onychophora of the World Diversity, Biology and Importance. Abstracts of 14th International Congress of Myriapodology. Staatliches Museum für Naturkunde, Görlitz. Peckiana 6: 123–124.
- Nefedieva JS, Nefediev PS, Sakhnevich MB, Dyachkov YuV (2014) Distribution of millipedes (Diplopoda) along an altitudinal gradient in the south of Lake Teletskoye, Altai Mts. In: Tuf IH, Tajovský K (Eds) 16th International Congress of Myriapodology. Book of Abstracts. Institute of Soil Biology, BC ASCR & Faculty of Science, Palacký University, Olomouc, 65.
- Özdikmen H (2008) New family and genus names, *Kirkayakidae* nom. nov. and *Kirkayakus* nom. nov., for the millipedes (Diplopoda: Chordeumatida). *Munis Entomology and Zoolology* 3(1): 342–344.

- Ryabinin NA, Ganin GN, Pankov AN (1988) [Effect of sulfuric acid industry wastes on soil invertebrate complexes]. *Ekologiya* 6: 29–37. [In Russian]
- Rybalov LB (2002) [Zonal-landscape change of soil invertebrate population in Yenisei Region of Middle Siberia and role of temperature adaptation in meridional (zonal) distribution of invertebrates]. *Russian Entomological Journal* 11(1): 77–86. [In Russian, with English summary]
- Shear WA (1988) Systematic position of the milliped species *Alloioopus solitaries* Attems and the genus *Ghilarovia* Gulička (Chordeumatida, Anthroleucosomatidae). *Myriapodologica* 2(8): 51–58.
- Shear WA (1990) On the Central and East Asian milliped family Diplomaragnidae (Diplopoda, Chordeumatida, Diplomaragnoidea). *American Museum Novitates* 2977: 1–40.
- Shelley RM, Sierwald P, Kiser SB, Golovatch SI (2000) Nomenclator generum et familiarum Diplopodorum II. A list of the genus and family-group names in the class Diplopoda from 1958 through 1999. Pensoft Publishers, Sofia-Moscow, 167 pp.
- StatSoft (2011) STATISTICA (data analysis software system), version 10. www.statsoft.com
- Stuxberg A (1876a) Myriopoder från Sibirien och Waigatsch on samlade under Nordenskiöldska expeditionen 1875. *Öfversigt k. Vetensk.-Akad. Förhandl.* 33(2): 11–38.
- Stuxberg A (1876b) On the Myriopoda, from Siberia and Waigatsch Island, collected during the expedition of Prof. Nordenskiöld, 1875. *Annals and Magazine of Natural History* 4(17): 306–318. doi: 10.1080/00222937608681955
- Vorobiova IG (1999) [An ecofaunistic characteristic of the myriapod population (Myriapoda) in a midflow region of the Yenisey River]. In: Biodiversity and life of soil system. Abstracts of XII All-Union Conference of Soil Zoology. Moscow, 33. [In Russian]
- Vorobiova IG, Rybalov LB, Rossolimo TE, Zalesskaya NT (2002) [Zonal and landscape distribution of the myriapod fauna and populations (Myriapoda) in the Yenisey River basin]. In: Study, conservation and reconstruction of ecosystem biodiversity in the Yenisey ecological transect: Wildlife, ethno-ecological investigations. Moscow 2: 60–71. [In Russian]